

On the dual equivalence of the Born-Infeld-Chern-Simons model coupled to dynamical U(1) charged matter

D. Bazeia,¹ A. Ilha,² J. R. S. Nascimento,¹ R. F. Ribeiro¹ and C. Wotzasek²

¹*Departamento de Física, Universidade Federal da Paraíba
58051-970 João Pessoa, Paraíba, Brasil*

²*Instituto de Física, Universidade Federal do Rio de Janeiro
21945-970, Rio de Janeiro, Brazil*

(February 1, 2008)

We study the equivalence between a nonlinear self-dual model (NSD) with the Born-Infeld-Chern-Simons (BICS) theory using an iterative gauge embedding procedure that produces the duality mapping, including the case where the NSD model is minimally coupled to dynamical, U(1) charged fermionic matter. The duality mapping introduces a current-current interaction term while at the same time the minimal coupling of the original nonlinear self-dual model is replaced by a non-minimal magnetic like coupling in the BICS side.

PACS numbers: 11.10.Kk, 11.10.Lm, 11.15.-q, 11.10.Cd

This work is devoted to the study of duality symmetry in the context of the Born-Infeld-Chern-Simons (BICS) theory [1], a generalization of the self-dual and Maxwell-Chern-Simons (MCS) duality, firstly shown by Deser and Jackiw [2] long time ago using the formalism of master Lagrangian. To this end we shall adopt a new gauge embedding formalism [3,4] that is alternative to the master Lagrangian approach.

Duality deals with the equivalence between models that describe the same physical phenomenon. This is a symmetry that is playing an important role in nowadays physics. It has received a spate of interest in recent research in diverse areas in field theory such as, supersymmetric gauge theories [5], string theories [6], sine-Gordon model [7], statistical systems [8] and, in the context of condensed matter models, applied for instance to planar high- T_C materials, Josephson junction arrays [9] and Quantum Hall Effect [10]. The existence of such a symmetry within a model may well have interesting consequences - it can be used to derive (exact) non perturbative results since swapping opposite coupling constant regimes allows a perturbative investigation of theories with large coupling constants. Moreover, it may be used to obtain information on the corresponding phase diagram, as it has been done in [8] where the relevant duality symmetry has been related to the modular group which, in that context, can be viewed as a generalization of the old Z_2 Kramers-Wannier duality of the 2-dimensional Ising model.

Another possibility, which is closely related to our interest here, is the odd-dimensional duality involving Chern-Simons term (CST) [11], whose paradigm is the equivalence between Self-Dual and Maxwell-Chern-Simons theories in (2+1) dimensions. This is a very special situation since the presence of the topological and gauge invariant Chern-Simons term is responsible for the essential features manifested by the three dimensional field theories, such as parity breaking and anomalous spin. It produces deep insights in unrelated areas in particle physics and condensed matter both from the theoretical and phenomenological points of view [12], as mentioned above. The high temperature asymptotic of four dimensional field theory models and the understanding of the universal behavior of the Hall conductance in interacting electron systems also stand as important illustrations of this topic.

Investigations of duality equivalence in D=3 involving CST has had a long history. In 1984, Deser and Jackiw [2] introduced the concept of master Lagrangian to prove that there are two different ways to describe the dynamics of a single, freely propagating spin one massive mode, either from the SD [13] or from the MCS theories [12]. In particular this result has been of great significance in order to extend the bosonization program from two to three dimensions with important phenomenological consequences [14].

On the other hand, the interest in the Born-Infeld theory [15] was recently revived in the investigations related to fluid mechanics and D-branes. For instance, both the relativistic Poincaré-invariant Born-Infeld model and the non-relativistic Galileo-invariant Chaplygin gas in $(d, 1)$ space-time dimensions descend from the very same reparametrization-invariant Nambu-Goto d-brane action in $(d + 1, 1)$ dimensions [16]. Also, in string theory it is well known that the Dp-brane is governed by an action S_{Dp} which is the sum of the Born-Infeld action with the Chern-Simons term, known as Born-Infeld-Chern-Simons (BICS) theory. For $p=2$ and 4, S_{Dp} can be derived from the actions of M2 and M5-brane in M-theory [17], and for $p=0$ it is obtained as the effective action of the Kaluza-Klein mode of the gravity multiplet in the eleven dimensional super-gravity. BICS theory is interesting in its own right since it gives the MCS theory as a limiting case. In this respect it becomes important to investigate the existence and to establish the dual equivalence with a generalized model having the SD model as a limiting case.

In this work we study the Chern-Simons duality for field theories including self-interaction and consider in particular the nonpolynomial BI action. To this end we apply an iterative gauge embedding procedure to construct an invariant

theory out of the nonlinear self-dual model leading to the BICS, including the coupling to dynamical matter fields. This is a gauge embedding procedure that changes the nature of the constraints of a given noninvariant theory. This is done with the inclusion of counter terms in the action, built with powers of the Euler vector which is enough to warrant the dynamical equivalence. Such construction discloses, in the language of constraints, hidden gauge symmetries in such systems. One can then consider the non-invariant model as the gauge fixed version of a gauge theory. The former reverts to the latter under certain gauge fixing conditions. By doing so we obtain a deeper and more illuminating interpretation of these systems. The associate gauge theory is therefore to be considered as the “gauge embedded” version of the original second-class theory. The advantage in having a gauge theory lies in the fact that the underlying gauge symmetry allows us to establish a chain of equivalence among different models by choosing different gauge fixing conditions.

The question we pose now is: how to find, using the gauge embedding approach for dual transformation, the (in principle, unknown) nonlinear SD theory that is dual to the BICS model (a gauge theory)? It should be clear that this is the reversed of the usual situation described before where the noninvariant model is known and one is looking for the (unknown) gauge embedded one. In any event, to answer this question we start by reviewing the (direct) method of gauge embedding [3,4] for the traditional SD model,

$$\mathcal{L}_{SD}^{(0)} = \frac{1}{2}m^2 f_\mu f^\mu - \frac{m}{2}\epsilon^{\mu\nu\rho} f_\mu \partial_\nu f_\rho, \quad (1)$$

where the superscript index is the iterative counter to be implemented here. Our basic goal is to transform the hidden symmetry of the Lagrangian (1) into a local gauge symmetry

$$\delta f_\mu = \partial_\mu \epsilon \quad (2)$$

with the lift of the global parameter ϵ into its local form, i.e.

$$\epsilon \rightarrow \epsilon(x, y, t) \quad (3)$$

The method works by looking for an (weakly) equivalent description of this theory which may be obtained by adding to the original Lagrangian (1) a function of the Euler vector K_μ

$$\mathcal{L}_{SD}^{(0)} \rightarrow \mathcal{L}_{SD}^{(0)} + f(K_\mu) \quad (4)$$

such that it vanishes on the space of solutions of (1), i.e., $f(0) = 0$. To find the specific form of this function that also induces a gauge symmetry into $\mathcal{L}_{SD}^{(0)}$ we work out iteratively. To this end we take the variation of $\mathcal{L}_{SD}^{(0)}$ to find the Euler vector as

$$K^\mu = m^2 f^\mu - m\epsilon^{\mu\nu\lambda} \partial_\nu f_\lambda \quad (5)$$

whose kernel gives the equations of motion of the model. Next we define a first-iterated Lagrangian as,

$$\mathcal{L}_{SD}^{(1)} = \mathcal{L}_{SD}^{(0)} - B_\mu K^\mu. \quad (6)$$

where the Euler vector has been imposed as a constraint with B_μ acting as a Lagrange multiplier.

The transformation properties of B_μ accompanying the basic field transformation (2) is chosen so as to cancel the variation of $\mathcal{L}_{SD}^{(0)}$, which gives

$$\delta B_\mu = \partial_\mu \epsilon \quad (7)$$

A simple algebra then shows

$$\begin{aligned} \delta \mathcal{L}_{SD}^{(1)} &= -B_\mu \delta K^\mu \\ &= -\delta \left(\frac{m^2}{2} B^2 \right) + m B_\mu \epsilon^{\mu\nu\lambda} \partial_\nu \delta f_\lambda \end{aligned} \quad (8)$$

where we have used (2) and (7). Because of (2), the second term in the r.h.s. of (8) vanishes identically leading to a second iterated Lagrangian,

$$\mathcal{L}_{SD}^{(2)} = \mathcal{L}_{SD}^{(1)} + \frac{m^2}{2} B^2 \quad (9)$$

that is gauge invariant under the combined local transformation of f_μ and B_μ .

We have therefore succeed in transforming the global SD theory into a locally invariant gauge theory. We may now take advantage of the Gaussian character of the auxiliary field B_μ to rewrite (9) as an effective action depending only on the original variable f_μ . To this end we use (9) to solve for the field B_μ (call this solution $\bar{B}_\mu$), and replace it back into (9) to find

$$\begin{aligned}\mathcal{L}_{eff} &= \mathcal{L}_{SD}^{(2)}|_{B_\mu=\bar{B}_\mu} = \mathcal{L}_{SD}^{(0)} - \frac{1}{2m^2}K^2 \\ &= \frac{1}{2}m^2 f_\mu f^\mu - \frac{m}{2}\varepsilon^{\mu\nu\rho} f_\mu \partial_\nu f_\rho \\ &\quad - \frac{1}{2m^2} [m^2 f^\mu - m\varepsilon^{\mu\nu\lambda} \partial_\nu f_\lambda]^2\end{aligned}\tag{10}$$

which can be rewritten to give the MCS theory,

$$\mathcal{L}_{MCS} = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} - \frac{m}{2}\varepsilon^{\mu\nu\rho}A_\mu\partial_\nu A_\rho.\tag{11}$$

It becomes clear from the above derivation that the difference between these two models is given by a function of the Euler vector of the SD model that vanishes over its space of solutions. This establishes the dynamical equivalence between the SD and the MCS theories.

Next, still in the direct situation, we consider the inclusion of self-interactions. To be specific let us consider a SD model with a quartic term,

$$\mathcal{L}_2 = \mathcal{L}_{SD} + \frac{\lambda}{4}(f_\mu f^\mu)^2.\tag{12}$$

The kinematics of the theory is controlled by the CST with the dynamics being determined by the potential that includes the mass term and the quartic self-interaction. Diagrammatically the SD model determines the basic propagator while the quartic term gives the vertex with strength λ . The basic idea of our approach is to reduce the self-interaction to a trilinear term. To this end we shall use the technique of auxiliary fields to reduce the interaction term to the form, $\sigma f_\mu f^\mu$, that may be treated as in the case above with the mass term being field dependent, plus a function of the auxiliary field. For the quartic term, only one auxiliary field is necessary,

$$\frac{\lambda}{4}(f_\mu f^\mu)^2 \rightarrow \frac{1}{2}\sigma f_\mu f^\mu + \frac{1}{\lambda}\sigma^2.\tag{13}$$

This operation has the effect of smooth out the quartic vertex very much like the old Fermi theory with the introduction of an intermediary field σ .

It is easy now to find the CS dual of this interacting SD model following the general prescription developed above. To this end we compute the Euler current K_μ relative to gauge transformations of the basic field f_μ keeping fixed the auxiliary field

$$K^\mu = (m^2 + \sigma) f^\mu - m\varepsilon^{\mu\nu\lambda} \partial_\nu f_\lambda.\tag{14}$$

The effective Lagrangian that results following the above procedure reads,

$$\begin{aligned}\mathcal{L}_{eff} &= \mathcal{L}_{SD} - \frac{1}{2m^2 + \sigma}K_\mu^2 + \frac{2}{\lambda}\sigma^2 \\ &= m\varepsilon^{\mu\nu\lambda}A_\mu\partial_\nu A_\lambda + \frac{2}{\lambda}\sigma^2 + \frac{F_{\mu\nu}F^{\mu\nu}}{2m^2 + \sigma}\end{aligned}\tag{15}$$

The final step is to solve the model for the field σ . The equation of motion gives

$$\sigma(\sigma + m^2)^2 = F_{\mu\nu}F^{\mu\nu}\tag{16}$$

Let us call $\bar{\sigma}$ the solution of this equation,

$$\bar{\sigma} = \bar{\sigma}(F_{\mu\nu}^2)\tag{17}$$

Bringing this solution back in the effective Lagrangian (15) gives the σ (Maxwell)-Chern-Simons theory that is dual to the quartic self-interacting SD theory.

$$\mathcal{L}_{\sigma CS} = m\epsilon^{\mu\nu\lambda}A_\mu\partial_\nu A_\lambda + \frac{2}{\lambda}\bar{\sigma}(F_{\mu\nu}^2) + \frac{F_{\mu\nu}^2}{2m^2 + \bar{\sigma}(F_{\mu\nu}^2)} \quad (18)$$

Of course a closed expression for $\bar{\sigma}$ in terms of the field tensor $F_{\mu\nu}$ is hard obtain in this case. There is however a nonlinear case where a closed expression is possible. This will be considered in the sequence.

We next consider the reversed situation mentioned above and discuss the duality equivalence involving the BICS theory. Let us say that some (yet) unknown nonlinear SD theory

$$\mathcal{L}_{NSD} = H(f_\mu f^\mu) - \frac{1}{2m}\epsilon^{\mu\nu\rho}f_\mu\partial_\nu f_\rho, \quad (19)$$

is the dual to the BICS with $H(f_\mu f^\mu)$ being nonlinear and possibly a nonpolynomial function of the $f_\mu f^\mu$. Our strategy is the assume that this term posses a linear representation, in terms of $f_\mu f^\mu$, given by an auxiliary field representation as

$$\mathcal{L}_{NSD} = g(\sigma) + \frac{1}{2}\sigma f_\mu f^\mu - \frac{1}{2m}\epsilon^{\mu\nu\rho}f_\mu\partial_\nu f_\rho. \quad (20)$$

The function $g(\sigma)$ is then determined such that the dual of (19) gives the BICS theory. To this end we repeat the steps above and compute the Euler vector,

$$K^\mu = \sigma f^\mu - \frac{1}{m}\epsilon^{\mu\nu\lambda}\partial_\nu f_\lambda \quad (21)$$

which allows us to find an effective action as

$$\begin{aligned} \mathcal{L}_{eff} &= \mathcal{L}_{NSD} - \frac{1}{2\sigma}K_\mu^2 \\ &= g(\sigma) + \frac{1}{2m}\epsilon^{\mu\nu\rho}A_\mu\partial_\nu A_\rho - \frac{1}{4m^2}\frac{1}{\sigma}F_{\mu\nu}F^{\mu\nu}. \end{aligned} \quad (22)$$

We have also relabeled $f_\mu \rightarrow A_\mu$ to reflect the inherent gauge invariance of the system. Next we solve for the field σ in terms of $F_{\mu\nu}^2$ to get,

$$\left[\sigma^2 \frac{d}{d\sigma} g(\sigma) \right]_{\sigma=\bar{\sigma}} + \frac{1}{4m^2}F_{\mu\nu}^2 = 0 \quad (23)$$

We must look at a conditional solution for this equation such that its substitution back into (22) should give the BI action,

$$g(\bar{\sigma}) - \frac{1}{4m^2}\frac{1}{\bar{\sigma}}F_{\mu\nu}^2 = \beta^2 \sqrt{1 - \frac{1}{2m^2\beta^2}F_{\mu\nu}^2} \quad (24)$$

The solution for these equations is given by

$$g(\sigma) = a\sigma + b\frac{1}{\sigma} \quad (25)$$

with the constants a and b being determined by (24) self-consistently as,

$$a = b = \frac{\beta^2}{2}. \quad (26)$$

Bringing this solution back into (22) gives

$$\mathcal{L}_{BICS} = \beta^2 \sqrt{1 - \frac{1}{2m^2\beta^2}F_{\mu\nu}^2} + \frac{1}{2m}\epsilon^{\mu\nu\rho}A_\mu\partial_\nu A_\rho. \quad (27)$$

which is the BICS theory we are looking for.

We are now in position to find the nonpolynomial function $H(f_\mu^2)$ in (19), that fixes the dual to the BICS theory, by bringing the solution (23)-(26) back into (20) and solving the equation of motion of the Lagrangian for the auxiliary field which, in this case, has a closed exact solution as,

$$\tilde{\sigma} = \frac{1}{\sqrt{1 + \frac{f_\mu f^\mu}{\beta^2}}} . \quad (28)$$

The auxiliary field is then eliminate from the theory by bringing it back into the Lagrangian (22) to produce,

$$\mathcal{L}_{NSD} = \beta^2 \sqrt{1 + \frac{f_\mu f^\mu}{\beta^2}} - \frac{1}{2m} \varepsilon^{\mu\nu\rho} f_\mu \partial_\nu f_\rho . \quad (29)$$

It is a simple algebra to show that indeed the equation of motion derived from $\mathcal{L}_{NSD}$ and $\mathcal{L}_{BICS}$ coincide. This solution has been found recently by Tripathy and Khare [1] by looking for a generalization of the equations of motion of the SD and MCS actions that have the correct limit for large values of β .

We are now in position to consider the minimal coupling of the NSD model with dynamical fermions. This generalizes the result of Gomes et al [18] into the BI context. Let us consider a Dirac field minimally coupled to a vector field specified by the NSD model [18], so that the Lagrangian becomes

$$\mathcal{L}_{min}^{(0)} = \mathcal{L}_{NSD} - e f_\mu J^\mu + \mathcal{L}_D , \quad (30)$$

where $J^\mu = \bar{\psi} \gamma^\mu \psi$, M is the fermion mass and the superscript index is the iterative counter. Here the Dirac Lagrangian is,

$$\mathcal{L}_D = \bar{\psi} (i \not{\partial} - M) \psi , \quad (31)$$

To implement the gauge embedding we follow the usual track and compute the Euler vector for the Lagrangian $\mathcal{L}_{NSD}$ given by (20), now in the presence of the fermionic current,

$$K^\mu = \sigma f^\mu - \frac{1}{m} \varepsilon^{\mu\nu\lambda} \partial_\nu f_\lambda - e J^\mu \quad (32)$$

The effective theory that comes out after the dualization procedure, with $f_\mu \rightarrow A_\mu$, is given by (22) with the Euler vector given by (32)

$$\begin{aligned} \mathcal{L}_{eff} = & g(\sigma) + \frac{1}{2m} \varepsilon^{\mu\nu\rho} A_\mu \partial_\nu A_\rho \\ & - \frac{1}{\sigma} \left[\frac{1}{4m^2} F_{\mu\nu}^2 + \frac{e^2}{2} J_\mu^2 + \frac{e}{m} \varepsilon^{\mu\nu\lambda} J_\mu \partial_\nu A_\lambda \right] \end{aligned} \quad (33)$$

with $g(\sigma)$ given by (25) and (26). Next the auxiliary field is eliminate in the usual way - from its equation of motion we find

$$\bar{\sigma} = \sqrt{1 - \frac{2}{\beta^2} \left(\frac{1}{4m^2} F_{\mu\nu}^2 + \frac{e^2}{2} J_\mu^2 + \frac{e}{m} \varepsilon^{\mu\nu\lambda} J_\mu \partial_\nu A_\lambda \right)} \quad (34)$$

which, when inserted back into (33), produces the dual BICS theory in the presence of dynamical fermions,

$$\begin{aligned} \mathcal{L}_{BICS} = & \frac{1}{2m} \varepsilon^{\mu\nu\rho} A_\mu \partial_\nu A_\rho \\ & + \beta^2 \sqrt{1 - \frac{2}{\beta^2} \left(\frac{1}{4m^2} F_{\mu\nu}^2 + \frac{e^2}{2} J_\mu^2 + \frac{e}{m} \varepsilon^{\mu\nu\lambda} J_\mu \partial_\nu A_\lambda \right)} . \end{aligned} \quad (35)$$

A simple inspection shows that the minimal coupling of the NSD model was replaced by a nonminimal magnetic Pauli type coupling and also the presence of a Thirring like current-current term. These features are characteristics of Chern-Simons dualities involving matter coupling [18,4]

ACKNOWLEDGMENTS: This work is partially supported by CNPq, PRONEX, CAPES, FAPERJ and FUJB, Brazilian Research Agencies. CW thanks the Physics Department of UFPB for the kind hospitality at the beginning of this investigation.

- [1] Prasanta K. Tripathy and Avinash Khare, Phys. Lett. B504 (2001) 152.
- [2] S. Deser and R. Jackiw, Phys. Lett. B 139 (1984) 2366.
- [3] A. Ilha and C. Wotzasek, Nucl. Phys. B, in press.
- [4] M. A. Anacleto, A. Ilha, J. R. S. Nascimento, R. F. Ribeiro and C. Wotzasek, Phys. Lett. B504 (2001) 268.
- [5] N. Seiberg and E. Witten, Nucl. Phys. B246 (1994) 19.
- [6] For a review see E. Kiritsis, Supersymmetry and Duality in Field Theory and String Theory, preprint hep-th/9911525 (1999).
- [7] D. Carpentier, J. Phys. A: Math. Gen. 32 (1999) 3865 and references therein.
- [8] J. L. Cardy, Nucl. Phys. B205 (1982) 17; J. L. Cardy and E. Rabinovici. Nucl. Phys. B205 (1982) 1.
- [9] M. C. Diamantini, P. Sodano and C. A. Trugenberger, Nucl. Phys. B448 (1995) 505; M. C. Diamantini, P. Sodano and C. A. Trugenberger, Nucl. Phys. B474 (1996) 641; for a general review see E. Fradkin, Field Theories of Condensed Matter systems, Addison-Wesley, Redwood City (1991).
- [10] Quantum Hall Effect, M. Stone, (ed.); Singapore: World Scientific (1992).
- [11] S. Chern and J. Simons, Ann. Math. 99 (1974) 49.
- [12] S. Schoenfeld, Nuc. Phys. B 185 (1981) 157; R. Jackiw and S. Templeton, Phys. Rev. D23 (1981) 2291; S. Deser, R. Jackiw and S. Templeton, Ann. Phys. 140 (1982) 372; Phys. Rev. Lett. 48 (1982) 372.
- [13] P. K. Townsend, K. Pilch and P. van Nieuwenhuizen, Phys. Lett. B 136 (1984) 38.
- [14] E. C. Marino, Phys. Lett. B263 (1991) 63; C. P. Burgess, C. A. Lutken and F. Quevedo, Phys. Lett. B336 (1994) 18; C. P. Burgess and F. Quevedo, Nucl. Phys. B421 (1994) 373; E. Fradkin and F. A. Schaposnik, Phys. Lett. B338 (1994) 253; R. Banerjee, Phys. Lett. B358 (1995) 297.
- [15] M. Born and M. Infeld, Proc. R. Soc. London A 144 (1934) 425.
- [16] R. Jackiw and A. P. Polychronakos, Commun. Math Phys. **207** (1999) 107; For a review see R. Jackiw, A Particle Field Theorist's Lectures on Supersymmetric, Non-Abelian Fluid Mechanics and d-Branes; physics/0010042.
- [17] S. P. de Alwis, Phys. Rev. D56 (1997) 7963; P. K. Townsend, Phys. Lett. B350 (1995) 184.
- [18] M. Gomes, L. C. Malacarne, A. J. da Silva, Phys. Lett. B439 (1998) 137.